

THE STRUCTURE OF THE FREE BOUNDARY IN THE FULLY NONLINEAR THIN OBSTACLE PROBLEM

XAVIER ROS-OTON AND JOAQUIM SERRA

ABSTRACT. We study the regularity of the free boundary in the fully nonlinear thin obstacle problem. Our main result establishes that the free boundary is C^1 near regular points.

1. INTRODUCTION

The aim of this paper is to study the regularity of free boundaries in thin obstacle problems.

1.1. Known results. The first regularity results for thin obstacle problems were already established in the seventies by Lewy [Lew68], Frehse [Fre77], Caffarelli [Caf79], and Kinderlehrer [Kin81]. In particular, for the Laplacian Δ , it was proved in [Caf79] that solutions are $C^{1,\alpha}$, for some small $\alpha > 0$.

The regularity of free boundaries, however, was an open problem during almost 30 years. One of the main difficulties in the understanding of free boundaries in thin obstacle problems is that there is not an a priori preferred order at which the solution detaches from the obstacle (blow-ups may have different homogeneities), as explained next.

In the classical (thick) obstacle problem it is not difficult to show that

$$0 < cr^2 \leq \sup_{B_r(x_0)} u \leq Cr^2 \quad (1.1)$$

at all free boundary points x_0 , where u is the solution of the problem (after subtracting the obstacle φ). Then, thanks to this, the blow-up sequence $u(x_0 + rx)/r^2$ converges to a global solution u_0 , and such solutions u_0 can be shown to be convex and completely classified; see [Caf98, Caf80] and [Caf77].

The situation is quite different in thin obstacle problems, in which one does not have (1.1). This was resolved for the first time in Athanasopoulos-Caffarelli-Salsa [ACS08], by using *Almgren's frequency function*. Thanks to this powerful tool, one

2010 *Mathematics Subject Classification.* 35R35; 35J60.

Key words and phrases. Thin obstacle problem; fully nonlinear; Signorini problem.

may take the blow-up sequence

$$\frac{u(x_0 + rx)}{\left(\int_{\partial B_r(x_0)} u^2 \right)^{1/2}},$$

and it converges to a *homogeneous* function u_0 of degree μ , for some $\mu > 1$. Then, by analyzing an eigenvalue problem on S^{n-1} , one can prove that

$$\mu < 2 \quad \implies \quad \mu = \frac{3}{2},$$

and for $\mu = \frac{3}{2}$ one can completely classify blow-ups. This leads to the optimal $C^{1, \frac{1}{2}}$ regularity of solutions and, using also a boundary Harnack inequality in “slit” domains, to the $C^{1, \alpha}$ regularity of the free boundary near *regular points* —those at which $\mu < 2$.

After the results of [ACS08], further regularity results for the free boundary have been obtained in [CSS08], [GP09], [GPS15], [DS14], [KPS15], [KRS15] and [BFR15].

1.2. Our setting. In this paper we study the fully nonlinear thin obstacle problem

$$\begin{cases} F(D^2u) \leq 0 & \text{in } B_1 \\ u \geq \varphi & \text{on } B_1 \cap \{x_n = 0\}, \\ F(D^2u) = 0 & \text{in } B_1 \setminus \{(x', 0) : u(x', 0) = \varphi(x')\}. \end{cases} \quad (1.2)$$

Here, $x = (x', x_n) \in \mathbb{R}^n$. When u is even with respect to the variable x_n , then the problem is equivalent to

$$\begin{aligned} F(D^2u) &= 0 & \text{in } B_1 \cap \{x_n > 0\} \\ \min(-u_{x_n}, u - \varphi) &= 0 & \text{on } B_1 \cap \{x_n = 0\}. \end{aligned} \quad (1.3)$$

Problem (1.3) was studied in [MS08], where Milakis and Silvestre proved that solutions u are $C^{1, \alpha}(B_{1/2})$ (for some small $\alpha > 0$) by following the ideas of [Caf79]. More recently, Fernández-Real extended the results of [MS08] to the general non-symmetric setting (1.2) in [Fer16].

Still, nothing was known about the regularity of the free boundary for this problem. The main difficulty in the study of such nonlinear thin free boundary problems is the lack of monotonicity formulas for fully nonlinear operators, which makes the proofs of [ACS08] non-applicable to the nonlinear setting.

1.3. Main results. We present here a new approach towards the regularity of thin free boundaries, and prove that for problem (1.2) the free boundary is C^1 near regular points.

As in [MS08, Fer16], we assume that the fully nonlinear operator F satisfies:

$$F \text{ is convex, with ellipticity constants } \lambda, \Lambda, \text{ and } F(0) = 0. \quad (1.4)$$

Our main result reads as follows.

Theorem 1.1. *Let F be as in (1.4). There exists $\epsilon_0 = \epsilon_0(\lambda, \Lambda) > 0$ for which the following holds.*

Let $u \in C(B_1)$ be any solution of (1.2), with $\varphi \in C^{1,1}$. Then, at each free boundary point $x_0 \in \partial\{u = \varphi\} \cap B_{1/2} \cap \{x_n = 0\}$ we have the following dichotomy:

$$(i) \text{ either } \sup_{B_r(x_0)} (u - \varphi) \geq c r^{2-\epsilon_0},$$

with $c > 0$,

$$(ii) \text{ or } \sup_{B_r(x_0)} (u - \varphi) \leq C_\epsilon r^{2-\epsilon} \quad \text{for all } \epsilon > 0.$$

Moreover, the set of points x_0 satisfying (i) is an open subset of the free boundary and it is locally a C^1 graph.

Notice that, for the Laplacian Δ , once we know that the free boundary is C^1 , then it can be proved that it is C^∞ ; see [DS14, KPS15] and also [RS15].

On the other hand, when F is the Laplacian Δ , at all free boundary points satisfying (i) the blow up is homogeneous of degree $3/2$, and thus all solutions are $C^{1, \frac{1}{2}}$. We do not expect this same exponent $3/2$ for all nonlinear operators $F(D^2u)$. A priori, each different operator F could have one (or more) different exponent μ , and thus in general solutions would be no better than $C^{1,\alpha}$ for some small $\alpha > 0$. Still, we show in Section 7 that

$$u \in C^{1, \frac{1}{2}-\delta}(B_{1/2}) \quad \text{whenever } |\Lambda - \lambda| \text{ is small enough;}$$

see Corollary 7.3.

We think that an interesting feature of our proof is that we establish the regularity of the free boundary without proving any homogeneity or uniqueness of blow-ups, a priori they could be non-homogeneous and/or non-unique. We do not classify blow-ups but only prove that they are 1D on $\{x_n = 0\}$, as explained next.

1.4. The proofs. To establish Theorem 1.1 we assume that x_0 is a regular free boundary point (i.e., (ii) does not hold at x_0), and do a blow-up. We have to do the blow-up along an appropriate subsequence, so that we get in the limit a global *convex* solution to (1.2), with zero obstacle, and with *subquadratic growth* at infinity. Then, we need to prove that blow-ups are 1D on $\{x_n = 0\}$, that is, the blow-up u_0 is a 1D function on $\{x_n = 0\}$, and in particular the contact set $\Omega^* = \{u_0 = 0\} \cap \{x_n = 0\}$ is a half-space.

To do this, we first notice that by a blow-down argument we may reduce to the case in which the convex set Ω^* is a convex cone Σ^* . Then, we separate into two cases, depending on the “size” of the convex cone Σ^* . If Σ^* has zero measure, then u_0 is in fact a global solution, and has subquadratic growth. By C^2 regularity estimates this is not possible, and thus Σ^* can not have zero measure. If Σ^* has nonempty interior, by convexity of u_0 this means that we have a cone of directional derivatives satisfying $\partial_\epsilon u_0 \geq 0$ in $\mathbb{R}^n$. Then, by a boundary Harnack type estimate (that we also establish here), we prove that all such derivatives have to be comparable in $\mathbb{R}^n$, and that this yields that the cone must be a half-space.

Once we have that blow-ups are 1D on $\{x_n = 0\}$, we show that the free boundary $\partial\{u = \varphi\}$ is Lipschitz in a neighborhood of any regular point x_0 , and C^1 at that point. Finally, by a barrier argument we show that the regular set is open —with all points in a neighborhood satisfying a uniform nondegeneracy condition. From here, we deduce that the free boundary is C^1 at every point in a neighborhood, with a uniform modulus of continuity.

Notice that an important step in the previous argument is the boundary Harnack type result for the derivatives $\partial_e u_0$, which solve an equation with bounded measurable coefficients in non-divergence form. The boundary Harnack principle for non-divergence equations is known to be false in $C^{0,\alpha}$ domains of $\mathbb{R}^n$ whenever $\alpha \leq \frac{1}{2}$; see [BB94]. Still, we prove here that a weaker version of the boundary Harnack principle holds in “slit” domains of the form $\mathbb{R}^n \setminus \Sigma^*$, where $\Sigma^* \subset \mathbb{R}^{n-1} \times \{0\}$ is a convex cone. The proof of such boundary Harnack type estimate is new, and we think it could be of independent interest.

Finally, notice also that our boundary Harnack type result allows us to show that blow-ups are 1D, but does *not* yield the $C^{1,\alpha}$ regularity of free boundaries. This is because the constants in such boundary Harnack estimate degenerate as the cone Σ^* contains two rays forming an angle approaching π .

1.5. Plan of the paper. The paper is organized as follows.

In Section 2 we construct some barriers that are needed in our proofs, and prove a maximum principle in $\mathbb{R}_+^n$ for functions u with sublinear growth. In Section 3 we establish our boundary Harnack type inequality for non-divergence equations with bounded measurable coefficients. In Section 4 we prove that global convex solutions with subquadratic growth to the fully nonlinear thin obstacle problem are necessarily 1D on $\{x_n = 0\}$. In Section 5 we show that at any regular free boundary point there is an appropriate rescaling such that the rescaled solutions converge in the C^1 norm to a global convex solution with subquadratic growth. In Section 6 we prove that the free boundary is flat Lipschitz by combining the results of Section 5 with a maximum principle argument. Finally, in Section 7 we show by a barrier argument that the regular set is open, which yields the C^1 regularity of the free boundary.

2. PRELIMINARIES AND TOOLS

We prove here some results that will be used in the paper. We will denote

$$M^+u = M^+(D^2u) \quad \text{and} \quad M^-u = M^-(D^2u),$$

the Pucci extremal operators; see [CC95] for their definition and basic properties.

Throughout the paper we call constants depending only on the dimension n and the ellipticity constants λ, Λ *universal constants*. Also, we denote B^+ the half ball $B \cap \{x_n > 0\}$, where B is some ball centered at some point on $\{x_n = 0\}$, and we denote by B^* , Σ^* , and Ω^* , “thin” balls, cones, and sets contained on $\{x_n = 0\}$.

2.1. Barriers. We first construct two barriers.

Lemma 2.1. *For $N = (n-1)\Lambda/\lambda$ the function*

$$\phi_0(x) = \begin{cases} \min\{1, |x'|^2 + N(2x_n - x_n^2)\} & \text{in } |x'| \leq 1, 0 \leq x_n \leq 1 \\ 1 & \text{elsewhere in } x_n \geq 0 \end{cases}$$

is continuous (viscosity) supersolution of $M^+\phi_0 \leq 0$ in $x_n > 0$.

Proof. We note that $|x'|^2 + N(2x_n - x_n^2) \geq |x'|^2 + |x_n|^2 \geq |x|$ and thus ϕ_0 is continuous. Also, where $\phi_0 < 1$ we have $M^+\phi_0 = 2(n-1)\Lambda - 2N\lambda \leq 0$. Thus, using that the minimum of two supersolutions is a supersolution we easily obtain that $M^+\phi_0 \leq 0$ in all of $\mathbb{R}^n$. $\square$

Lemma 2.2. *Let $a_i \geq 0$ with $\sum_{i=0}^{\infty} a_i < \infty$. Then, the function*

$$\phi(x) = \sum_{i=0}^k 2^i a_i \phi_0(2^{-i}x)$$

is a continuous (viscosity) supersolution of $M^+\phi \leq 0$ in all of $x_n > 0$. Moreover, ϕ satisfies

$$2^j a_j \leq \phi \quad \text{in } \overline{B_{2^{j+1}}^+} \setminus \overline{B_{2^j}^+} \quad (2.1)$$

and

$$\phi \leq C \left(\sum_{i=0}^j 2^i a_i + \sum_{i=j}^{\infty} a_i \right) \quad \text{in } \overline{B_{2^j}^+} \quad (2.2)$$

where C is a universal constant.

Proof. Let ϕ_0 be the supersolution from Lemma 2.1. We then consider, for $k \geq 0$

$$\phi^k(x) = \sum_{i=0}^k 2^i a_i \phi_0(2^{-i}x)$$

On one hand, we have

$$M^+\phi^k(x) \leq \sum_{i=0}^k 2^{i-2} a_i M^+\phi_0(2^{-i}x) \leq 0.$$

On the other hand, whenever $k \geq j$ and $|x| \geq 2^j$ we have

$$\phi^k(x) \geq 2^j a_j \phi_0(2^{-j}x) \geq 2^j a_j, \quad (2.3)$$

since we readily check that $\phi_0 \geq \min\{1, |x'|^2 + |x_n|^2\} = 1$ outside B_1^+ (in $x_n > 0$).

Finally, we note that $\phi_0 \leq C \min\{1, |x'| + |x_n|\}$ and thus

$$\phi^k(x) \leq C \sum_{i=0}^k 2^i a_i \min\{1, 2^{-i}|x|\} \leq C \left(\sum_{i=0}^j 2^i a_i + \sum_{i=j}^{\infty} a_i \right) \quad \text{for } x \in B_{2^j}^+. \quad (2.4)$$

Then, the monotone increasing sequence ϕ^k converges locally uniformly in $\{x_n > 0\}$ to some function $\phi = \phi^\infty$. By the stability of viscosity supersolutions under uniform convergence we have $M^+\phi \leq 0$ in all of $\mathbb{R}^n$. That ϕ satisfies the other conditions of the lemma is easily verified letting $k \rightarrow \infty$ in (2.3) and (2.4). $\square$

The following subsolution will be used in the proof of our boundary Harnack inequality.

Lemma 2.3. *Given $\rho \in (0, 1)$ and a ball $B^* = B_r^*(z)$ ($z \in \mathbb{R}^{n-1}$), with B^* contained in B_1^* , there is a function $\phi \in C(B_1)$ satisfying*

$$\begin{cases} M^-\phi \geq \chi_{B_{1-\rho}} & \text{in } B_1 \setminus B^* \\ \phi \geq 0 & \text{in } B_1 \\ \phi \leq C\chi_{B^*} & \text{on } B_1^* \\ \phi = 0 & \text{on } \partial B_1 \end{cases} \quad (2.5)$$

where C depends only on ρ , B^* and universal constants.

Proof. Let g_0 be the restriction to ∂B_1^+ of the function $\max\{0, 1 - (x - z)^2/r^2\}$ and $f_0(x) = f_0(|x|)$ be a radial nonincreasing function with $f_0 = 0$ for $|x| \geq 1 - \rho/2$ and $f_0 = 1$ for $|x| \leq 1 - \rho$.

For $\kappa \in (0, 1)$ small, we let ψ be the solution to

$$\begin{cases} M^-\psi_\kappa = \kappa f_0 & \text{in } B_1^+ \\ \psi = g_0 & \text{on } \partial B_1^+ \end{cases} \quad (2.6)$$

Let us show that κ small enough (depending only on ρ and B^*) we have $\psi \geq 0$ in B_1^+ .

Indeed, by the strong maximum principle and Hopf's lemma, for $\kappa = 0$ we have

$$\psi_0 \geq \delta_0 > 0 \quad \text{in } B_{1-\rho/4} \cap \{x_n > \rho/4\}.$$

Thus, by the uniqueness of solution to (2.6) and the stability of viscosity solutions we deduce that

$$\psi_\kappa \geq \delta_0/2 > 0 \quad \text{in } B_{1-\rho/4} \cap \{|x_n| > \rho/4\}. \quad (2.7)$$

for κ small.

Next, for N large enough the function $\eta = \exp(-N|x|) - \exp(-N\rho/2)$ satisfies

$$M^-\eta = \left(\lambda N^2 - \frac{\Lambda N(n-1)}{|x|} \right) \eta > 0 \quad \text{in } \{|x| \geq \rho/4\} \cap \{\eta > 0\}. \quad (2.8)$$

Thus, we have $M^-\eta \geq c > 0$ in $\{\rho/4 \leq |x| \leq \rho/2\}$ and using $\frac{\delta_0}{2}\eta(x - x_0)$ as a barrier (by below) with x_0 on $\{|x'| \leq 1 - \rho/2, x_n = \rho/2\}$, and by (2.7) we obtain

$$\psi_\kappa \geq 0 \quad \text{in } B_{1-\rho/2}^+ \quad (2.9)$$

when κ is chosen small enough.

Finally, from (2.9) it follows that (still for κ small) we have $\psi_\kappa \geq 0$ in all of B_1^+ . Here we are using that $f_0 = 0$ in the half annulus $B_1 \setminus B_{1-\delta/2}$.

To end the proof, we let ϕ be the even reflection of the previous $\frac{1}{\kappa}\psi_\kappa$ with respect to the variable x_n multiplied by a large positive constant C . Then, using that ϕ will have a negative wedge on $B_1^* \setminus B^*$ it not difficult to verify that it will satisfy all the requirements of the lemma. $\square$

2.2. A maximum principle in $\mathbb{R}_+^n$ and construction of 1D solutions. We next prove the following.

Lemma 2.4. *Let u satisfy*

$$\sup_{B_1^+} |u| + \sum_{i=0}^{\infty} 2^{-i} \sup_{B_{2^{i+1}}^+ \setminus B_{2^i}^+} |u| < \infty \quad (2.10)$$

and

$$\begin{cases} M^-u \leq 0 & (\text{resp. } M^+u \geq 0) & \text{in } \{x_n > 0\} \\ u \geq 0 & (\text{resp. } u \leq 0) & \text{on } \{x_n = 0\}. \end{cases}$$

Then, $u \geq 0$ (resp. $u \leq 0$) in $\{x_n > 0\}$.

For this, we need the following.

Lemma 2.5. *Let (a_k) be a sequence such that $a_k \geq 0$ and $\sum_{k \geq 1} a_k < \infty$. Then, there exists a sequence (b_k) such that $b_k/a_k \geq 1$, $\lim b_k/a_k = \infty$, and $\sum_{k \geq 1} b_k < \infty$.*

Proof. The result is probably well known, we give here a proof for completeness.

Let us define $s_k = \sum_{j \geq k} a_j$. Note that may (and do) assume that $s_1 = 1$. Let

$$b_k = \frac{a_k}{\sqrt{\sum_{j \geq k} a_j}} = \frac{a_k}{\sqrt{s_k}} \geq a_k.$$

Notice that $\lim b_k/a_k = \infty$, since $s_k \rightarrow 0$. Then, we have

$$b_k = \frac{s_k - s_{k+1}}{\sqrt{s_k}} \leq 2\sqrt{s_k} - 2\sqrt{s_{k+1}}, \quad (2.11)$$

where we used that $2\sqrt{x} - 2\sqrt{y} \geq (x - y)/\sqrt{x}$ for all $x \geq y$ (this follows from the mean value theorem). Therefore, by (2.11), we find

$$\sum_{k \geq 1} b_k \leq 2\sqrt{s_1} < \infty,$$

and the lemma is proved. $\square$

We now give the:

Proof of Lemma 2.4. Let $a_i := 2^{-i} \sup_{B_{2^{i+1}}^+ \setminus B_{2^i}^+} |u|$. By assumption $\sum a_i < \infty$ and then, by Lemma 2.5, there exists b_i increasing such that $1 \leq b_i/a_i \rightarrow \infty$ and $\sum b_i < \infty$. Then, we consider

$$\phi(x) := -\sup_{B_1^+} |u| - \sum_{i=0}^{\infty} 2^i b_i \phi_0(2^{-i}x),$$

where ϕ_0 is the supersolution in the proof of Lemma 2.2. Exactly as in the proof of Lemma 2.2 we find that ϕ is subsolution in all of $\{x_n > 0\}$. Then, using that $u \geq 0$ on $\{x_n = 0\}$, that $b_i/a_i \rightarrow \infty$, and the maximum principle, we obtain $u \geq -\epsilon\phi$ in all of $\{x_n \geq 0\}$ for every $\epsilon > 0$. Thus $u \geq 0$ in all of $\{x_n \geq 0\}$. $\square$

As a consequence of Lemma 2.4, we find the following.

Proposition 2.6 (Extensions). *Given $g : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$ continuous satisfying*

$$\sup_{B_1^*} |g| + \sum_{i=0}^{\infty} 2^{-i} \sup_{B_{2^{i+1}}^* \setminus B_{2^i}^*} |g| < \infty$$

there exist a unique function u belonging to $C(\{x_n > 0\})$ which satisfies (2.10) and

$$\begin{cases} M^+ u = 0 & \text{in } \{x_n > 0\} \\ u = g & \text{on } \{x_n = 0\}. \end{cases}$$

We then denote $E^+ g := u$.

Similarly $E^- g := -E^+(-g)$ is the unique solution, among functions satisfying (2.10), of the previous problem with M^+ replaced by M^- .

Proof. Let $a_i = 2^{-i} \sup_{B_{2^{i+1}}^* \setminus B_{2^i}^*} |g|$ and

$$\phi(x) := \sup_{B_1^*} |g| + \sum_{i=0}^{\infty} 2^i a_i \phi_0(2^{-i} x),$$

By Lemma 2.2 we have $\phi \geq g$ in $x_n = 0$ and $M^+ \phi \leq 0$ in $x_n > 0$. On the other hand, using (2.2) we find

$$\begin{aligned} \sum_{j=0}^{\infty} 2^{-j} \sup_{B_{2^j}} \phi &\leq \sum_{j=0}^{\infty} 2^{-j} \left(\sup_{B_1^*} |g| + C \sum_{i=0}^j 2^{i-j} a_i + C \sum_{j=0}^{\infty} \sum_{i=j}^{\infty} 2^{-j} a_i \right) \\ &\leq 2 \sup_{B_1^*} |g| + C \sum_{i=0}^{\infty} \sum_{j=i}^{\infty} 2^{i-j} a_i + 2C \sum_{i=0}^{\infty} a_i \\ &\leq 2 \sup_{B_1^*} |g| + 4C \sum_{i=0}^{\infty} a_i < \infty. \end{aligned}$$

Thus in particular ϕ satisfies (2.10) with u replaced by ϕ .

Now we note that ϕ and $-\phi$ are respectively a supersolution and a subsolution of the problem $M^+ u = 0$ in $\{x_n > 0\}$, $u = g$ on $\{x_n = 0\}$. Then, we can prove the existence of a continuous viscosity solution between $-\phi$ and ϕ in several standard ways.

One option is to choose any continuous extension $\bar{g}$ of g to $\{x_n > 0\}$ such that $|\bar{g}| \leq \phi$ and to solve in large balls $M^+ u_R = 0$ in B_R^+ , $u = \bar{g}$ in ∂B_R^+ . Letting $R \uparrow \infty$ and using the stability of viscosity solutions under local uniform converge, we find a solution of the of the problem in all of $x_n > 0$. The barriers $\pm\phi$ guarantee the

convergence. Another option is to proof the existence of a solution in the half space directly by Perron's method.

The uniqueness of viscosity solution to this problem among continuous functions u satisfying (2.10) is a straightforward consequence of the maximum principle in Lemma 2.4 and the fact that the difference w of two solutions satisfies $M^+w \geq 0$ and $M^-w \leq 0$ in $\{x_n > 0\}$, and $w = 0$ on $\{x_n = 0\}$. $\square$

We next construct 1D solutions in $\mathbb{R}_+^2$.

Proposition 2.7. *For any $\beta \in (0, 1)$, let us consider the function $\varphi_\beta^\pm(x, y) := E^\pm(x_+)^beta$ in $\mathbb{R}_+^2$. Then,*

(a) *We have*

$$\partial_y \varphi_\beta^+ = \overline{C}(\beta) x^{\beta-1} \quad \text{in } \{x > 0\} \cap \{y = 0\},$$

$$\partial_y \varphi_\beta^- = \underline{C}(\beta) x^{\beta-1} \quad \text{in } \{x > 0\} \cap \{y = 0\}.$$

The constants $\overline{C}$ and $\underline{C}$ depend only on β and ellipticity constants.

(b) *The functions $\overline{C}(\beta)$ and $\underline{C}(\beta)$ are continuous in β , and there are*

$$0 < \beta_1 < \frac{1}{2} < \beta_2 < 1$$

such that

$$\overline{C}(\beta_1) = 0 \quad \text{and} \quad \underline{C}(\beta_2) = 0.$$

Moreover, β_1 and β_2 are unique.

(c) *For any small $\delta > 0$, we have*

$$\frac{1}{2} - \delta < \beta_1 < \frac{1}{2} < \beta_2 < \frac{1}{2} + \delta \quad \text{whenever} \quad |\Lambda - 1| + |\lambda - 1| \leq \delta/C,$$

with C universal.

We will need the following auxiliary result.

Lemma 2.8. *Let $w_k = E^+g_k$ (resp. $w_k = E^-g_k$) where*

$$\sum_{i \geq 1} 2^{-i} \sup_{B_{2^i}^*} |g_k| \leq C, \tag{2.12}$$

and

$$\|g_k\|_{C^{1,\alpha}(\overline{B_{1/2}^*})} \leq C, \tag{2.13}$$

for some $\alpha \in (0, 1)$, with C independent of k .

Suppose that, for some $g \in C(\mathbb{R}^{n-1} \times \{0\})$

$$\sum_{i \geq 1} 2^{-i} \sup_{B_{2^i}} |g_k - g| \rightarrow 0 \quad \text{on } \{x_n = 0\}. \tag{2.14}$$

Then, $|\partial_{x_n} w_k - \partial_{x_n} w|(0) \rightarrow 0$, where $w = E^+g$ (resp. $w = E^-g$).

Proof. We first show that $w_k \in C^{1,\alpha}(\overline{B_{1/4}^+})$, with a bound independent of k , and that $w_k \rightarrow w$ uniformly in $\overline{B_{1/4}^+}$.

Indeed, it follows from (2.12) and from Lemma 2.2 (see also the proof of Proposition 2.6) that $\|w_k\|_{L^\infty(B_1^+)} \leq C$, with C independent of k . Then, by the $C^{1,\alpha}$ estimates up to the boundary (see [CC95]) using (2.13) we obtain that $\|w_k\|_{C^{1,\alpha}(\overline{B_{1/4}^+})} \leq C$.

On the other hand, $w_k - w$ is a viscosity solution of $M^-(w_k - w) \leq 0 \leq M^+(w_k - w)$ in $\{x_n > 0\}$. Then by (2.14) —using again Lemma 2.2— we find $\sup_{B_1^+}(w_k - w) \rightarrow 0$.

Since all the w_k are uniformly $C^{1,\alpha}(\overline{B_{1/4}^+})$ and converge uniformly to w in $\overline{B_1^+}$ we find in particular $w_k \rightarrow w$ in $C^1(\overline{B_{1/4}^+})$. Thus, $|\partial_{x_n} w_k - \partial_{x_n} w|(0) \rightarrow 0$. $\square$

We now give the:

Proof of Proposition 2.7. (a) It follows by the scaling properties of $M^\pm$ and by uniqueness of $E^\pm$ that $\varphi_\beta^\pm$ are homogeneous functions of degree β . Thus, part (a) follows, with

$$\overline{C}(\beta) = \partial_y \varphi_\beta^+(1, 0), \quad \underline{C}(\beta) = \partial_y \varphi_\beta^-(1, 0).$$

(b) It follows from Lemma 2.8 —translating the origin to the point $(1, 0)$ — that $\partial_y \varphi_\beta^\pm(1, 0) \rightarrow \partial_y \varphi_\beta^\pm(1, 0)$. As a consequence, $\overline{C}(\beta)$ and $\underline{C}(\beta)$ are continuous in $\beta \in (0, 1)$. Although for $\beta = 0$, the function $(x_+)^beta = \chi_{x>0}$ has a discontinuity, we can easily adapt the proof of Lemma 2.8 to this situation by using that the only discontinuity point is at $(0, 0)$ and that the solution is bounded near this discontinuity point.

Note instead that a similar continuity property is not true as $\beta \uparrow 1$, since we approach the critical growth and hence we can not guarantee that $\|\varphi_\beta^\pm\|_{L^\infty(\overline{B_{1/4}^+(1,0)})}$ stays bounded as $\beta \uparrow 1$. In fact, we will show later on in this proof that this L^∞ norm diverges.

Now, when $\beta = 0$, as said above $\varphi_\beta^\pm(x, 0) = \chi_{\{x>0\}}$ and Hopf lemma implies that $\partial_y \varphi_\beta^\pm(1, 0) < 0$. Thus,

$$\lim_{\beta \downarrow 0} \underline{C}(\beta) \leq \lim_{\beta \downarrow 0} \overline{C}(\beta) < 0.$$

On the other hand, we claim that

$$\overline{C}(\beta) \geq \underline{C}(\beta) \geq \frac{c}{1-\beta} \rightarrow \infty \quad \text{as } \beta \uparrow 1. \quad (2.15)$$

Indeed, let ψ be the subsolution of Lemma 2.3, with $r_0 = \frac{1}{4}$ and extended by zero outside B_1 . Consider the new subsolution

$$\psi_k(x, y) = \sum_{i=0}^k 2^{\beta i} \psi(2^{-i}x - \frac{1}{2}, 2^{-i}y),$$

which satisfies $M^-\psi_k \geq 0$ in all of $\{y > 0\}$.

Note that, since $r_0 = 1/4$, the functions we have $\psi(2^{-i}x - 1/2, 2^{-i}y)$ have disjoint supports at $y = 0$. Thus, we find

$$\psi_k(x, 0) \leq 2^{i\beta} \chi_{\{0 < x < 2^i\}} \quad \text{for all } k \text{ and } i$$

In particular $2^{-\beta}\psi_k \leq (x_+)^{\beta}$ on $\{y = 0\}$. Now, for fixed β we readily show, using Lemma 2.4 and Proposition 2.6, that

$$2^{-\beta}\psi_k \leq \varphi_{\beta}^{-} = E^{-}(x_+)^{\beta} \quad (\text{for all } k). \quad (2.16)$$

But note that, by Lemma 2.3, at $x = \frac{1}{4}$ we have $\psi_k(\frac{1}{4}, 0) = 0$ and thus

$$\psi_k\left(\frac{1}{4}, y\right) = \sum_{i=0}^k 2^{(\beta-1)i} (\partial_i \psi) \left(2^{-i} - \frac{1}{2}, 2^{-i}y\right) \geq c \frac{1 - 2^{(\beta-1)k}}{1 - 2^{\beta-1}} y.$$

for $|y| < 1/2$. Letting $k \rightarrow \infty$, using (2.16), and recalling that φ_{β}^{-} is homogeneous of degree β we obtain

$$\varphi_{\beta}^{-}(x, y) \geq \frac{c}{1 - 2^{\beta-1}} y \geq \frac{c}{1 - \beta} y \quad \text{for } x \in \left(\frac{1}{2}, \frac{3}{2}\right), \quad y \in (0, 1)$$

for some $c > 0$ universal.

As $\beta \uparrow 1$, thus $\varphi_{\beta}^{-}(x, y)$ is a nonnegative solution in $Q = (1, 2, 3/2) \times (0, 1)$ with trace x^{β} on $(1, 2, 3/2) \times \{y = 0\}$ and that is arbitrarily large in $(1, 2, 3/2) \times (1/2, 1)$. Then it is immediate to show that there is a quadratic polynomial P satisfying $M^{-}P \geq 0$ (subsolution), such that P touches φ_{β}^{-} by below in $\bar{Q}$ at the point $x = 1$, $y = 0$, and with $\partial_y P(1, 0)$ arbitrarily large. Thus $\underline{C}(\beta)$ is arbitrarily large as $\beta \rightarrow 1$ —with a growth $c/(1 - \beta)$ —, finishing the proof of the claim (2.15).

Finally, as said before, $\bar{C}$ and $\underline{C}$ are continuous functions. Thus, there are $0 < \beta_1 \leq \beta_2 < 1$ such that $\bar{C}(\beta_1) = 0$ and $\underline{C}(\beta_2) = 0$.

The uniqueness of the exponents β_1 and β_2 follows by a simple contact argument. Indeed, if $\beta < \beta'$ then some translation (to the right) of the function φ_{β}^{+} touches $\varphi_{\beta'}^{+}$ by below at some point on $\{x > 0, y = 0\}$. But since the two functions are homogeneous the sign of their vertical derivatives is the same on all of $\{x > 0, y = 0\}$. This shows that $\text{sign}(\bar{C}(\beta')) > \text{sign}(\bar{C}(\beta))$, where the strict inequality is a consequence of Hopf Lemma. This implies that the zero of $\bar{C}$ is unique. The same argument applies to $\underline{C}$.

Finally, using the same contact argument to compare $\varphi_{\beta_1}^{+}$ and $\varphi_{\beta_2}^{-}$ with the harmonic extension of $(x_+)^{1/2}$ (i.e. the solution for the Laplacian), we obtain $\beta_1 < \frac{1}{2} < \beta_2$.

(c) Let ψ be the solution of

$$\psi(x, 0) = (x_+)^{\frac{1}{2}-\delta} \quad \text{on } \{y = 0\},$$

$$\Delta\psi = -\kappa r^{-\frac{3}{2}-\delta} \quad \text{in } \{y > 0\},$$

where $r = \sqrt{x^2 + y^2}$. Notice that ψ is homogeneous of degree $\frac{1}{2} - \delta$ in $\mathbb{R}^2$.

Notice also that when $\kappa = 0$ then $\psi_y(x, 0) = -c(\delta)x^{-\frac{1}{2}-\delta} < 0$ for $x > 0$. Thus, if κ is small, we will have $\psi_y(x, 0) \leq -\frac{1}{2}c(\delta)x^{-\frac{1}{2}-\delta}$ for $x > 0$. In fact, a simple computation shows that $c(\delta) \geq c\delta$ for δ small. Thus, by linearity, we may take $\kappa \geq c\delta > 0$, too.

Let us now check that, if $|\Lambda - 1| + |\lambda - 1| \leq \gamma$, with $\gamma > 0$ small, then

$$M^+\psi \leq 0 \quad \text{in} \quad \{y > 0\}.$$

For this, notice that by homogeneity of ψ we only need to check it on ∂B_1 , where ψ is C^2 . Also, notice that

$$M^+\psi = \lambda\Delta\psi + (\Lambda - \lambda)(\text{sum of positive eigenvalues of } D^2\psi),$$

so that

$$M^+\psi \leq \lambda\Delta\psi + C(\Lambda - \lambda) \leq -\lambda\kappa + C\gamma \leq -c\delta + C\gamma \leq 0$$

provided that $\gamma \leq \delta/C$.

Thus,

$$M^+\psi \leq 0 \quad \text{on} \quad \{y > 0\},$$

$$\psi_y \leq 0 \quad \text{on} \quad \{y = 0, x > 0\},$$

$$\psi \text{ is homogeneous of degree } \frac{1}{2} - \delta.$$

This, and the same contact argument as before, yields $\frac{1}{2} - \delta < \beta_1$. Repeating the same argument with $\frac{1}{2} + \delta$, we get $\frac{1}{2} + \delta > \beta_2$, and thus the proposition is proved. $\square$

As a consequence, we have the following.

Corollary 2.9. *Given $e \in S^{n-2}$, let*

$$w_0^+(x) := \varphi_{\beta_1}^+(x' \cdot e, |x_n|), \quad w_0^-(x) := \varphi_{\beta_2}^-(x' \cdot e, |x_n|),$$

where $\varphi_{\beta}^{\pm}$ and β_1, β_2 are given by Proposition 2.7. Then,

$$\begin{cases} M^{\pm}w_0^{\pm} = 0 & \text{in } \mathbb{R}^n \setminus (\{x' \cdot e \leq 0\} \cap \{x_n = 0\}) \\ w_0^{\pm} = 0 & \text{on } \{x' \cdot e \leq 0\} \cap \{x_n = 0\}. \end{cases}$$

The functions w_0^+ and w_0^- are homogeneous of degree β_1 and β_2 , respectively, and $0 < \beta_1 < \frac{1}{2} < \beta_2 < 1$.

Moreover, $\frac{1}{2} - \delta < \beta_1 < \frac{1}{2} < \beta_2 < \frac{1}{2} + \delta$ whenever $|\Lambda - 1| + |\lambda - 1| \leq \delta/C$.

Proof. The result follows from Proposition 2.7, and taking into account that since $M^{\pm}w_0^{\pm} = 0$ in $\{x_n \neq 0\}$ and $w_0^{\pm}$ are C^1 at points on $\{x' \cdot e > 0\} \cap \{x_n = 0\}$, then they also solve the equation therein. $\square$

2.3. A maximum principle type Lemma. We finally prove the following Lemma, similar to [ACS08, Lemma 5].

Lemma 2.10. *Let c_0, c_1 be given positive constants with $c_1 < \sqrt{\lambda/(9n\Lambda)}$ —i.e. universally small enough. Then, there exists $\sigma > 0$ for which the following holds.*

Assume $v \in C(\overline{B_1})$ satisfies

- $M^-v \leq \sigma$ in $B_1 \setminus \Omega^*$, with $\Omega^* \subset \{x_n = 0\}$
- $v = 0$ on Ω^*
- $v \geq c_0 > 0$ for $|x_n| \geq c_1 > 0$
- $v \geq -\sigma$ in B_1

Then, $v \geq 0$ in $B_{1/2}$. Moreover, $v \geq c_2|x_n|$ in $B_{1/2}$, for some $c_2 > 0$ (small).

Proof. Let us prove that $v \geq 0$ in $B_{1/2}$. Once this is proved, then $v \geq c_2|x_n|$ follows from the standard subsolution of Hopf's lemma —see (2.8)— provided that σ is small enough.

Assume there is $z = (z', z_n) \in B_{1/2} \cap \{|x_n| < c_1\}$ such that $v(z) < 0$. Let

$$Q = \left\{ (x', x_n) : |x' - z'| \leq \frac{1}{3}, |x_n| \leq c_1 \right\}$$

and

$$P(x) = |x' - z'|^2 - \frac{n\Lambda}{\lambda}x_n^2.$$

Notice that $M^+P = -\Lambda$.

Define

$$w = v + \delta P,$$

where $\delta > 0$ is such that $0 < C\sigma < \delta < c_0/C$, with C large enough. Then, we have

- $w(z) = v(z) - \delta\Lambda z_n^2 < 0$
- $M^-w \leq M^-v + \delta M^+P \leq \sigma - \delta\Lambda \leq 0$ outside Ω^*
- $w \geq 0$ on Ω^*

Thus, w must have a negative minimum on ∂Q .

On $\partial Q \cap \{|x_n| = c_1\}$ we have

$$w \geq c_0 - \delta \frac{n\Lambda}{\lambda} c_1^2 \geq 0.$$

On $\partial Q \cap \{|x' - z'| = 1/3\} \cap \{0 \leq |x_n| \leq c_1\}$, we have $v \geq -\sigma$, so that

$$w \geq -\sigma + \delta \left(\frac{1}{9} - \frac{n\Lambda}{\lambda} c_1^2 \right) \geq 0.$$

Hence, $w \geq 0$ on ∂Q and we have reached a contradiction. Therefore, $v \geq 0$ in $B_{1/2}$, as desired. $\square$

3. A BOUNDARY HARNACK INEQUALITY

We prove here a boundary Harnack inequality in “slit” cones, for solutions that are monotone in some “outwards” directions. More precisely, we establish the following.

Proposition 3.1. *Let $\Sigma^* \subset \mathbb{R}^{n-1} \times \{0\}$ be some nonempty closed convex cone satisfying*

$$\Sigma^* \subset \left\{ \frac{x}{|x|} \cdot e \leq -\varepsilon \right\} \quad (3.1)$$

for some $e \in S^{n-2}$ and $\varepsilon \in (0, 1/8)$. Let θ_1, θ_2 be unit vectors in $\mathbb{R}^{n-1} \times \{0\}$ with $-\theta_i \in \Sigma^*$.

Assume that $u_1, u_2 \in C(B_1)$ satisfy

$$M^+(au_1 + bu_2) \geq 0 \quad \text{in } B_1 \setminus \Sigma^* \quad (3.2)$$

for all $a, b \in \mathbb{R}$,

$$u_1 = u_2 = 0 \quad \text{on } B_1^* \cap \Sigma^*.$$

Assume also $u_i \geq 0$ in B_1^+ , $\sup_{B_{\varepsilon/2}} u_1 = \sup_{B_{\varepsilon/2}} u_2$, and u_i is monotone nondecreasing in the direction θ_i in all of B_1 —that is, $u_i(\bar{x}) \geq u_i(x)$ whenever $\bar{x} - x = t\theta_i$ for some $t \geq 0$ and $x, \bar{x} \in B_1$.

Then,

$$\frac{1}{C\varepsilon^{-M}} u_2 \leq u_1 \leq C\varepsilon^{-M} u_2 \quad \text{in } \overline{B_{\varepsilon/4}},$$

where C and M are positive universal constants.

Proof. We may and do assume that

$$\sup_{B_{\varepsilon/2}} u_i = 1. \quad (3.3)$$

Step 1. We define

$$A_\varepsilon := B_{7/8} \cap \{x \cdot e \geq \varepsilon/4\}.$$

We first prove that that

$$0 < C_\varepsilon^{-1} \leq \inf_{A_\varepsilon} u_i \leq 1 \quad (3.4)$$

where $C_\varepsilon := C\varepsilon^{-M}$ for some positive universal constants C and M . Throughout the proof C_ε denotes a constant of this form though C and M may vary from line to line.

Indeed, first note that by taking the four choices $a = \pm 1, b = 0$ and $a = 0, b = \pm 1$ in (3.10) we obtain that u_i are viscosity solutions of

$$M^- u_i \leq 0 \leq M^+ u_i \quad \text{in } B_1 \setminus \Sigma^*.$$

Thus, using a standard chain of interior Harnack inequalities we have

$$\sup_{A_\varepsilon} u_i \leq C_\varepsilon \inf_{A_\varepsilon} u_i.$$

On the other hand, let us show that

$$\text{given } x \in B_{\varepsilon/2} \text{ exist } \bar{x} \in A_\varepsilon, t \geq 0 \text{ such that } \bar{x} - x = t\theta_i$$

Indeed, if $x \in B_{\varepsilon/2}$ we have $x \cdot e > -\varepsilon/2$ and thus, using (3.9) the point $\bar{x} = x + \frac{3}{4}\theta_i$ satisfies

$$\bar{x} \cdot e \geq -\varepsilon/2 + 3\varepsilon/4 \geq \varepsilon/4.$$

Here we have used that $\theta_i \cdot e \geq \varepsilon$ since $-\theta_i$ are unit vectors in Σ^* and we have (3.9). In addition, $\bar{x} \in B_{7/8}$ since $|\frac{3}{4}\theta_i| = 3/4$ and $|x| = \varepsilon/2 \leq 1/8$.

Thus, using the monotonicity of u_i in the direction θ_i we have that

$$1 = \sup_{B_{\varepsilon/2}} u_i \leq \sup_{A_\varepsilon} u_i \leq C_\varepsilon \inf_{A_\varepsilon} u_i \leq C_\varepsilon \sup_{B_{\varepsilon/2}} u_i = C_\varepsilon,$$

where for the last inequality we have used that $A_\varepsilon \cap B_{\varepsilon/2} \neq \emptyset$.

Thus, (3.4) follows.

Step 2. We next prove that, with C_ε as above,

$$u_1 \geq C_\varepsilon^{-1} u_2 \quad \text{in } B_{\varepsilon/4}^*. \quad (3.5)$$

We consider the rescaled solutions $\bar{u}_i(x) = u_i(\frac{\varepsilon}{2}x)$. Then, $\bar{u}_1, \bar{u}_2 \in C(B_1)$ satisfy

$$M^+(a\bar{u}_1 + b\bar{u}_2) \geq 0 \quad \text{in } B_2 \setminus \Sigma^* \quad (3.6)$$

for all $a, b \in \mathbb{R}$, and

$$\bar{u}_1 = \bar{u}_2 = 0 \quad \text{on } B_2^* \cap \Sigma^*.$$

In addition we have $\bar{u}_i \geq 0$ in B_2 . $\sup_{B_1} \bar{u}_i = 1$ —recall (3.3)—, and, by Step 1,

$$C_\varepsilon^{-1} \leq \inf_{B_1 \cap \{e \cdot x \geq 1/4\}} \bar{u}_i.$$

Using again a chain of interior Harnack inequalities we obtain

$$C_\varepsilon^{-1} \leq \inf_{B^*} \bar{u}_i, \quad (3.7)$$

where $B^* = B_{1/4}^*(z)$ for $z = e/2$.

Fix $\rho = 1/10$. Let $\eta \in C^2(\overline{B_1})$ be some smooth “cutoff” function with $\eta = 1$ for $|x| \geq 1 - \rho$ and $\eta = 0$ in $B_{1/2}$. Let us call

$$C_1 := \sup_{B_1} M^+ \eta = \sup_{B_{1-\rho}} M^+ \eta$$

Let ϕ be the subsolution of Lemma 2.3—with $\rho = 1/10$ and $B^* = B_{1/4}^*(z)$ for $z = e/2$, as before.

We will show next that, for $C_\varepsilon \geq 1$ large enough,

$$C_\varepsilon \bar{u}_1 + \eta \geq \bar{u}_2 + C_1 \phi \quad \text{in } B_1. \quad (3.8)$$

Indeed, on the one hand since $0 \leq \bar{u}_i \leq 1$ in B_1 we and $\eta = 1$ for $|x| \geq 1 - \rho$ we have and $\phi = 0$ on ∂B_1 we have that (7.5) holds on ∂B_1 . On the other hand we have

$$M^-(C\bar{u}_1 + \eta - \bar{u}_2 - C_1 \phi) \leq M^+ \eta - C_1 M^- \phi \leq C_1 \chi_{B_{1-\rho}} - C_1 \chi_{B_{1-\rho}} \leq 0 \quad \text{in } B_1 \setminus B^*$$

while, using (3.7)

$$C_\varepsilon \bar{u}_1 + \eta - \bar{u}_2 - C_1 \phi \geq (C_\varepsilon \bar{u}_1 - \bar{u}_2) + (C_\varepsilon \bar{u}_1 - C_1 \phi) \geq 0 \quad \text{in } B^*$$

where we recall that C is a constant of the type $C\varepsilon^{-M}$ with C and M universal and varying from line to line.

Thus, (7.5) follows using by the maximum principle. Finally, since $\phi \geq 0$ and $\eta = 0$ in $B_{1/2}$ from (7.5) we deduce that

$$C_\varepsilon \bar{u}_1 \geq \bar{u}_2 \quad \text{in } B_{1/2}$$

and thus after rescaling we obtain (3.5).

Finally, since the roles of $\bar{u}_1$ and $\bar{u}_2$ are interchangeable we obtain the comparability of $\bar{u}_1$ and $\bar{u}_2$ in $\overline{B_{1/8}^+}$. Rescaling back, we obtain that u_1 and u_2 are comparable in $B_{\varepsilon/4}$, as desired. $\square$

As a consequence we obtain the following.

Corollary 3.2. *Let $\Sigma^* \subset \mathbb{R}^{n-1} \times \{0\}$ be some nonempty closed convex cone satisfying*

$$\Sigma^* \subset \left\{ \frac{x}{|x|} \cdot e \leq -\varepsilon \right\} \quad (3.9)$$

for some $e \in S^{n-2}$ and $\varepsilon \in (0, 1/8)$. Let θ_1, θ_2 be unit vectors in $\mathbb{R}^{n-1} \times \{0\}$ with $-\theta_i \in \Sigma^*$.

Assume that $u_1, u_2 \in C(\mathbb{R}^n)$ satisfy

$$M^+(au_1 + bu_2) \geq 0 \quad \text{in } \mathbb{R}^n \setminus \Sigma^* \quad (3.10)$$

for all $a, b \in \mathbb{R}$,

$$u_1 = u_2 = 0 \quad \text{on } \Sigma^*.$$

Assume also $u_i \geq 0$ in $\mathbb{R}^n$, $\sup_{B_1} u_1 = \sup_{B_1} u_2$, and u_i is monotone nondecreasing in the direction θ_i in all of $\mathbb{R}^n$ —that is, $u_i(\bar{x}) \geq u_i(x)$ whenever $\bar{x} - x = t\theta_i$ for some $t \geq 0$.

Then,

$$\frac{1}{C\varepsilon^{-M}} u_2 \leq u_1 \leq C\varepsilon^{-M} u_2 \quad \text{in all of } \mathbb{R}^n.$$

where C and M are positive universal constants.

Proof. We may assume that $\sup_{B_{1/2}} u_1 = \sup_{B_{1/2}} u_2 = 1$.

Let $R \geq 4$ arbitrary. Consider the two rescaled functions $\bar{u}_1$ and $\bar{u}_2$ defined by

$$\bar{u}_i(x) = \frac{u_i(Rx)}{C_i} \quad \text{for } C_i = \|u_i\|_{L^\infty(B_R)}.$$

By Proposition 3.1 we obtain that

$$C_\varepsilon^{-1} \bar{u}_2 \leq \bar{u}_1 \leq C_\varepsilon \bar{u}_2 \quad \text{in } \overline{B_{1/8}},$$

where $C_\varepsilon = C\varepsilon^{-M}$ with C and M universal constants.

Thus, using that

$$1 = \|u_i\|_{L^\infty(B_{1/2})} = C_i \|\bar{u}_i\|_{L^\infty(B_{1/(2R)})}$$

Since we have that $\|\bar{u}_1\|_{L^\infty(B_{1/(2R)})}$ and $\|\bar{u}_2\|_{L^\infty(B_{1/(2R)})}$ are comparable (recall that $R \geq 4$) we obtain that C_1 and C_2 are comparable and thus, scaling back, that

$$C_\varepsilon^{-1}u_2 \leq u_1 \leq C_\varepsilon u_2 \quad \text{in } \overline{B_{R/8}}.$$

Since R can be taken arbitrarily large the Corollary follows. $\square$

4. GLOBAL SOLUTIONS

In this Section we prove that any global solution to the obstacle problem with subquadratic growth must be 1D on $\{x_n = 0\}$.

Theorem 4.1. *Let F be as in (1.4), and $u \in C(\mathbb{R}^n)$ be any viscosity solution of*

$$\begin{cases} F(D^2u) \leq 0 & \text{in } \mathbb{R}^n \\ F(D^2u) = 0 & \text{in } \mathbb{R}^n \setminus \Omega^* \\ u = 0 & \text{on } \Omega^* \\ u \geq 0 & \text{on } \{x_n = 0\}, \end{cases} \quad (4.1)$$

with

$$u(0) = 0, \quad \nabla u(0) = 0. \quad (4.2)$$

Assume that u satisfies the following growth control

$$\|u\|_{L^\infty(B_R)} \leq R^{2-\epsilon} \quad \text{for all } R \geq 1. \quad (4.3)$$

Then, either $u \equiv 0$, or

$$u(x) = u_0(e \cdot x', x_n) \quad \text{and} \quad \{u(x', 0) = 0\} = \{e \cdot x' \leq 0\}$$

for some $e \in S^{n-2}$. Moreover, u_0 is convex in the x' variables.

We will need the following intermediate steps in the proof of Theorem 4.1.

Lemma 4.2. *Let F be as in (1.4), and $u \in C(\mathbb{R}^n)$ be any viscosity solution of*

$$F(D^2u) = 0 \quad \text{in } \mathbb{R}^n,$$

with $u(0) = 0$ and $\nabla u(0) = 0$. Assume that u satisfies the growth control (4.3). Then, $u \equiv 0$.

Proof. By interior $C^{1,1}$ estimates [CC95]—here we use the convexity of the operator—we have

$$\|D^2u\|_{L^\infty(B_1)} \leq C.$$

Applying the same estimate to the rescaled function $u(Rx)/R^{2-\epsilon}$, we find

$$\|D^2u\|_{L^\infty(B_R)} \leq CR^{-\epsilon},$$

for any $R \geq 1$. Letting $R \rightarrow \infty$, we deduce that u is affine. Since $u(0) = 0$ and $\nabla u(0) = 0$, it must be $u \equiv 0$. $\square$

We next prove the following.

Proposition 4.3. *Let F be as in (1.4), and $u \in C(\mathbb{R}^n)$ be any viscosity solution of (4.1)-(4.2)-(4.3) which is convex in the $x' = (x_1, \dots, x_{n-1})$ variables.*

Assume in addition that $\Sigma^* = \{u = 0\} \cap \{x_n = 0\}$ is a closed convex cone with nonempty interior and vertex at the origin. Then, either $u \equiv 0$ or

$$\Sigma^* = \{x' \cdot e \leq 0\}$$

for some $e \in S^{n-2}$.

Proof. Assume that u is not identically zero and that Σ^* is not a half-space.

Notice that if Σ^* contains a line $\{te' : t \in \mathbb{R}\}$ then by convexity of u we will have $u(x + te') = u(x)$ for all $t \in \mathbb{R}$, $x \in \mathbb{R}^n$. Hence, if Σ^* contains a line, u is a solution in dimension $n - 1$. Therefore, by reducing the dimension n if necessary, we may assume that Σ^* contains no lines.

In particular,

$$\Sigma^* \subset \left\{ \frac{x'}{|x'|} \cdot e \leq -\varepsilon \right\} \quad (4.4)$$

for some $e \in S^{n-2}$ and some $\varepsilon > 0$.

Let $\varepsilon > 0$ be the largest positive number for which (4.4) holds. Let $e_1 \in S^{n-2}$ be such that $-e_1 \in \Sigma^*$ and $-e_1 \cdot e = -\varepsilon$.

Since $-e \in \Sigma^*$ and $-e_1 \in \Sigma^*$, then by convexity of u we have

$$w = \partial_e u \geq 0 \quad \text{and} \quad w_1 = \partial_{e_1} u \geq 0 \quad \text{on} \quad \{x_n = 0\}.$$

Moreover, since Σ^* contains no lines, then these two functions are positive in $\{x_n = 0\} \setminus \Sigma^*$. Moreover, we have

$$M^+(aw + bw_1) \geq 0 \quad \text{in} \quad \mathbb{R}^n \setminus \Sigma^*$$

for all $a, b \in \mathbb{R}$. Furthermore, the convexity of u and the growth control (4.3) yield

$$\|w\|_{L^\infty(B_R)} + \|w_1\|_{L^\infty(B_R)} \leq CR^{1-\epsilon}.$$

By the maximum principle in Lemma 2.4, this implies

$$w = \partial_e u \geq 0 \quad \text{and} \quad w_1 = \partial_{e_1} u \geq 0 \quad \text{in} \quad \mathbb{R}^n.$$

Therefore, by the boundary Harnack type principle in Corollary 3.2, this means that

$$\partial_{e_1} u \geq c \partial_e u \quad \text{in} \quad \mathbb{R}^n.$$

Equivalently, $\partial_{e_1 - ce} u \geq 0$. But then this yields $-(e_1 - ce) \in \Sigma^*$, which combined with $-(e_1 - ce) \cdot e = -\varepsilon - c$ is a contradiction with (4.4). $\square$

Using Lemma 4.2 and Proposition 4.3, we can now give the:

Proof of Theorem 4.1. If $u \equiv 0$ there is nothing to prove. By the (local) semi-convexity estimates in [Fer16] applied (rescaled) to a sequence of balls with radius converging to infinity, we readily prove u is convex in the x' variables. Thus, Ω^* is convex.

If $\Omega^* = \{x' \cdot e \leq 0\}$ for some $e \in S^{n-2}$, then by convexity we have $u(x', 0) = u_0(x' \cdot e, 0)$, and thus $u(x) = u_0(x' \cdot e, x_n)$, where u_0 is a 2D solution to the problem.

We next prove that if Ω^* is not a half-space, then there is no solution u .

Assume by contradiction that Ω^* is not a half-space and that u is a nonzero solution. Then, we do a blow-down argument, as follows.

For $R \geq 1$ define

$$\theta(R) = \sup_{R' \geq R} \frac{\|u\|_{L^\infty(B_{R'})}}{(R')^{2-\epsilon}}.$$

Note that $0 < \theta(R) < \infty$ and that it is nonincreasing.

For all $m \in \mathbb{N}$ there is $R'_m \geq m$ such that

$$(R'_m)^{\epsilon-2} \|u_m\|_{L^\infty(B_R)} \geq \frac{\theta(m)}{2} \geq \frac{\theta(R'_m)}{2}.$$

Then the blow down sequence

$$u_m(x) := \frac{u(R'_m x)}{(R'_m)^{2-\epsilon} \theta(R'_m)}$$

satisfies the growth control

$$\|u_m\|_{L^\infty(B_R)} \leq R^{2-\epsilon} \quad \text{for all } R \geq 1$$

and also

$$\|u_m\|_{L^\infty(B_1)} \geq \frac{1}{2}.$$

By $C^{1,\alpha}$ estimates [Fer16] and the Arzelà-Ascoli theorem, the sequence u_m converges (up to a subsequence) locally uniformly in C^1 to a function u_∞ satisfying

$$\|u_\infty\|_{L^\infty(B_R)} \leq R^{2-\epsilon} \quad \text{for all } R \geq 1, \tag{4.5}$$

$$\|u_\infty\|_{L^\infty(B_1)} \geq \frac{1}{2}, \tag{4.6}$$

and

$$\begin{cases} F(D^2 u_\infty) = 0 & \text{in } \mathbb{R}^n \setminus \Sigma^* \\ F(D^2 u_\infty) \leq 0 & \text{in } \mathbb{R}^n \\ D^2 u_\infty \geq 0 & \text{in } \mathbb{R}^n \\ u_\infty = 0 & \text{in } \Sigma^*, \end{cases} \tag{4.7}$$

where Σ^* is the blow-down of the convex set Ω^* . Notice that, by convexity, since Ω^* was not a half-space, then Σ^* is not a half-space.

If Σ^* has nonempty interior, by Proposition 4.3 there is no solution u . If Σ^* has empty interior, then by $C^{1,\alpha}$ regularity of u we get $u_{x_n} = 0$ in all of $\{x_n = 0\}$. But using Lemma 4.2, this yields $u \equiv 0$ as well.

Thus, if Ω^* is not a half-space there is no nonzero solution u , as claimed. $\square$

We also prove the following.

Corollary 4.4. *Let F be as in (1.4), and $\beta_1 \in (0, \frac{1}{2})$ be given by Corollary 2.9. Let $u \in C(\mathbb{R}^n)$ be any viscosity solution of (4.1) satisfying (4.2) and*

$$\|u\|_{L^\infty(B_R)} \leq R^{1+\beta} \quad \text{for all } R \geq 1, \quad (4.8)$$

with $\beta < \beta_1$. Then, $u \equiv 0$.

Proof. By Theorem 4.1, we know that $u(x) = u_0(x' \cdot e, x_n)$, with u_0 convex in the first variable and vanishing on $\{x_1 \leq 0\} \cap \{x_2 = 0\}$. Thus, we only need to prove the result in dimension $n = 2$. We denote $v = \partial_{x_1} u \geq 0$ in $\mathbb{R}^2$. Notice that

$$\begin{cases} M^+v \geq 0, & M^-v \leq 0 & \text{in } \mathbb{R}^2 \setminus (\{x_1 \leq 0\} \cap \{x_2 = 0\}) \\ v = 0 & & \text{on } \{x_1 \leq 0\} \cap \{x_2 = 0\}. \end{cases}$$

Notice also that, by convexity and (4.8), we have $\|v\|_{L^\infty(B_R)} \leq CR^\beta$.

We now use the supersolution given by Corollary 2.9. Indeed, let $w = w_0^+$ be the homogeneous function of degree β_1 satisfying

$$\begin{cases} M^+w = 0 & \text{in } \mathbb{R}^2 \setminus (\{x_1 \leq 0\} \cap \{x_2 = 0\}) \\ w = 0 & \text{on } \{x_1 \leq 0\} \cap \{x_2 = 0\}. \end{cases}$$

Then, using interior Harnack inequality, a simple application of the maximum principle yields

$$0 \leq v \leq Cw \quad \text{in } B_2 \setminus B_1.$$

Here, we used that $\|v\|_{L^\infty(B_3)} \leq C$. By comparison principle, we deduce

$$0 \leq v \leq Cw \quad \text{in } B_2.$$

Repeating the same argument at all scales $R \geq 1$ —using the rescaled functions $R^{-\beta_1}w(Rx) = w(x)$ and $R^{-\beta_1}v(Rx)$ —, we find

$$0 \leq v \leq CR^{\beta-\beta_1}w \quad \text{in } B_{2R} \setminus B_R.$$

Here, we used that $\|v\|_{L^\infty(B_{3R})} \leq CR^\beta$.

By comparison principle, the previous inequality yields

$$0 \leq v \leq CR^{\beta-\beta_1}w \quad \text{in } B_R,$$

and thus letting $R \rightarrow \infty$ we find $v \equiv 0$. This means that $u(x_1, x_2) = \psi(x_2)$, for some function ψ . But since $F(D^2u) = 0$ in $\{x_2 > 0\}$ and in $\{x_2 < 0\}$, then $u(x_1, x_2) = ax_2$, and since $\nabla u(0) = 0$, then $u \equiv 0$, as desired. $\square$

5. REGULAR POINTS AND BLOW-UPS

We start in this section the study of free boundary points. For this, we use some ideas from [CRS16].

After a translation, we may assume that the free boundary point is located at the origin. Moreover, by subtracting a plane, we may assume that

$$u(0) = 0 \quad \text{and} \quad \nabla u(0) = 0.$$

Moreover, we assume

$$\|u\|_{L^\infty(B_1)} = 1, \quad \|\varphi\|_{C^{1,1}} \leq 1.$$

We say that a free boundary point is regular whenever (ii) in Theorem 1.1 does *not* hold, that is:

Definition 5.1. We say that $0 \in \partial\{u = \varphi\}$ is a *regular* free boundary point if

$$\limsup_{r \downarrow 0} \frac{\|u\|_{L^\infty(B_r)}}{r^{2-\epsilon}} = \infty$$

for some $\epsilon > 0$. We say that it is a regular point with exponent ϵ and modulus ν if

$$\sup_{\rho \leq r \leq 1} \frac{\|u\|_{L^\infty(B_r)}}{r^{2-\epsilon}} \geq \nu(\rho)$$

where $\nu(\rho)$ is a given nonincreasing function satisfying $\nu(\rho) \rightarrow \infty$ as $\rho \downarrow 0$.

The main result of this section is the following.

Proposition 5.2. *Assume that 0 is a regular free boundary point with exponent ϵ and modulus ν . Then, given $\delta > 0$, there is $r > 0$ such that the rescaled function*

$$v(x) := \frac{u(rx)}{\|u\|_{L^\infty(B_r)}}$$

satisfies

$$|v - u_0| + |\nabla v - \nabla u_0| \leq \delta \quad \text{in } B_1, \quad (5.1)$$

for some global convex solution u_0 of (4.1)-(4.2)-(4.3), with $\|u_0\|_{L^\infty(B_1)} = 1$. The constant r depends only on δ, ϵ, ν, n , and λ, Λ .

To prove this, we need the following intermediate step.

Lemma 5.3. *Given $\delta > 0$, there is $\eta = \eta(\delta, \epsilon, n, \lambda, \Lambda) > 0$ such that the following statement holds.*

Let φ be such that $\|\varphi\|_{C^{1,1}} \leq \eta$, and let $v \geq 0$ be a function satisfying $v(0) = 0$, $\nabla v(0) = 0$,

$$\begin{aligned} F(D^2v) &= 0 & \text{in } B_{1/\eta} \setminus \{x_n = 0\} \\ \min(-F(D^2v), v - \varphi) &= 0 & \text{on } B_{1/\eta} \cap \{x_n = 0\}, \end{aligned} \quad (5.2)$$

and

$$\|v\|_{L^\infty(B_1)} = 1, \quad \|v\|_{L^\infty(B_R)} \leq CR^{2-\epsilon} \quad \text{for } 1 \leq R \leq 1/\eta. \quad (5.3)$$

Then,

$$|v - u_0| + |\nabla v - \nabla u_0| \leq \delta \quad \text{in } B_1,$$

for some global convex solution u_0 of (4.1)-(4.2)-(4.3), with $\|u_0\|_{L^\infty(B_1)} = 1$.

Proof. The proof is by a compactness. Assume by contradiction that for some $\delta > 0$ we have a sequence $\eta_k \rightarrow 0$, fully nonlinear convex operators F_k with ellipticity constants λ, Λ , obstacles φ_k with $\|\varphi_k\|_{C^{1,1}} \leq \eta_k$, and functions $v_k \geq 0$ satisfying $v_k(0) = 0$, $\nabla v_k(0) = 0$, (5.2), and (5.3), but such that

$$\|v_k - u_0\|_{C^1(B_1)} \geq \delta \quad \text{for all global solution } u_0 \quad \text{with} \quad \|u_0\|_{L^\infty(B_1)} = 1. \quad (5.4)$$

By the estimates in [Fer16, MS08], we have that v_k are $C^{1,\alpha}$ in B_R , $R < 1/\eta_k$, with an estimate

$$\|v_k\|_{C^{1,\alpha}(B_R)} \leq C(R) \quad \text{for all} \quad 1 \leq R \leq 1/2\eta_k.$$

Thus, up to taking a subsequence, the operators F_k converge (locally uniformly as Lipchitz functions of the Hessian) to some fully nonlinear convex operator F with ellipticity constants λ, Λ . Likewise, the functions v_k converge in $C_{\text{loc}}^1(\mathbb{R}^n)$ to a function v_∞ , which by stability of viscosity solutions —see [CC95]— is a global convex solution to the obstacle problem (4.1) and satisfying (4.2) and (4.3).

By the classification result Theorem 4.1, we have

$$v_\infty \equiv u_0, \quad \text{for some global solution } u_0.$$

Moreover, by (5.3) we have

$$\|u_0\|_{L^\infty(B_1)} = \|v_\infty\|_{L^\infty(B_1)} = 1.$$

We have shown that $v_k \rightarrow u_0$ in the C^1 norm, uniformly on compact sets. In particular, (5.4) is contradicted for large k , and thus the lemma is proved. $\square$

To prove Proposition 5.2 we will also need the following.

Lemma 5.4. *Assume $w \in L^\infty(B_1)$ satisfies $\|w\|_{L^\infty(B_1)} = 1$, and*

$$\sup_{\rho \leq r \leq 1} \frac{\|w\|_{L^\infty(B_r)}}{r^{2-\epsilon}} \geq \nu(\rho) \rightarrow \infty \quad \text{as } \rho \rightarrow 0.$$

Then, there is a sequence $r_k \downarrow 0$ for which $\|w\|_{L^\infty(B_{r_k})} \geq \frac{1}{2}r_k^\mu$, and for which the rescaled functions

$$w_k(x) = \frac{w(r_k x)}{\|w\|_{L^\infty(B_{r_k})}}$$

satisfy

$$|w_k(x)| \leq C(1 + |x|^\mu) \quad \text{in } B_{1/r_k},$$

with $C = 2$. Moreover, we have

$$0 < 1/k \leq r_k \leq (\nu(1/k))^{-1/\mu}.$$

Proof. Let

$$\theta(\rho) := \sup_{\rho \leq r \leq 1} r^{-\mu} \|w\|_{L^\infty(B_r)}.$$

By assumption, we have

$$\theta(\rho) \geq \nu(\rho) \rightarrow \infty \quad \text{as } \rho \downarrow 0.$$

Note that θ is nonincreasing.

Then, for every $k \in \mathbb{N}$ there is $r_k \geq \frac{1}{k}$ such that

$$(r_k)^{-\mu} \|w\|_{L^\infty(B_{r_k})} \geq \frac{1}{2} \theta(1/k) \geq \frac{1}{2} \theta(r_k). \quad (5.5)$$

Note that since $\|w\|_{L^\infty(B_1)} = 1$ then

$$(r_k)^{-\mu} \geq \frac{1}{2} \theta(1/k) \geq \frac{1}{2} \nu(1/k),$$

and hence

$$0 < 1/k \leq r_k \leq (\nu(1/k))^{-1/\mu}.$$

Moreover, we have $\theta(r_k) \geq 1$, and thus $\|w\|_{L^\infty(B_{r_k})} \geq \frac{1}{2} r_k^\mu$.

Finally, by definition of θ and by (5.5), for any $1 \leq R \leq 1/r_k$ we have

$$\|w_k\|_{L^\infty(B_R)} = \frac{\|w\|_{L^\infty(B_{r_k R})}}{\|w\|_{L^\infty(B_{r_k})}} \leq \frac{\theta(r_k R)(r_k R)^\mu}{\frac{1}{2}(r_k)^\mu \theta(r_k)} \leq 2R^\mu.$$

In the last inequality we used the monotonicity of θ . $\square$

We now give the:

Proof of Proposition 5.2. Let $r_k \rightarrow 0$ be the sequence given by Lemma 5.4 (with $\mu = 2 - \epsilon$). Then, the functions

$$u_k(x) = \frac{u(r_k x)}{\|u\|_{L^\infty(B_{r_k})}}$$

satisfy

$$|u_k(x)| \leq C(1 + |x|^\mu) \quad \text{in } B_{1/r_k},$$

and

$$\|u_k\|_{L^\infty(B_1)} = 1, \quad u_k(0) = 0, \quad \nabla u_k(0) = 0. \quad (5.6)$$

Moreover, they are solutions to the obstacle problem in B_{1/r_k} , i.e.,

$$\begin{aligned} F(D^2 u_k) &= 0 & \text{in } B_{1/r_k} \setminus \{x_n = 0\} \\ \min(-F(D^2 u_k), u_k - \varphi_k) &= 0 & \text{on } B_{1/r_k} \cap \{x_n = 0\}, \end{aligned}$$

where

$$\|\varphi_k\|_{C^{1,1}} = \frac{\|\varphi(r_k \cdot)\|_{C^{1,1}}}{\|u_k\|_{L^\infty(B_{r_k})}} \leq \frac{C(r_k)^2}{(r_k)^{2-\epsilon}} = C(r_k)^\epsilon$$

converges to 0 uniformly as $k \rightarrow \infty$. Therefore, by Lemma 5.3 for k large enough (so that $(r_k)^\epsilon \leq (\nu(1/k))^{-1/(2-\epsilon)} \leq \eta$) we have

$$|v - u_0| + |\nabla v - \nabla u_0| \leq \delta \quad \text{in } B_1,$$

for some global convex solution u_0 of (4.1)-(4.2)-(4.3), with $\|u_0\|_{L^\infty(B_1)} = 1$, as desired. $\square$

6. LIPSCHITZ REGULARITY OF THE FREE BOUNDARY

We now prove that the free boundary is Lipschitz in a neighborhood of any regular point x_0 .

Proposition 6.1. *Assume that 0 is a regular free boundary point with exponent ϵ and modulus ν . Then, there exists $e \in S^{n-1} \cap \{x_n = 0\}$ such that for any $\ell > 0$ there exists $r > 0$ for which*

$$\partial_\tau u \geq 0 \quad \text{in } B_r \quad \text{for all } \tau \cdot e \geq \frac{\ell}{\sqrt{1+\ell^2}}, \quad \tau \in S^{n-1} \cap \{x_n = 0\}.$$

In particular, the free boundary is Lipschitz in B_r , with Lipschitz constant ℓ . The constant r depends only on $\ell, \epsilon, \nu, n, \lambda, \Lambda$.

To prove this, we need the following.

Lemma 6.2. *Let $u_0(x) = u_0(x' \cdot e, x_n)$ be a global solution of (4.1)-(4.2)-(4.3), with $\|u_0\|_{L^\infty(B_1)} = 1$. Let $\tau \in S^{n-1} \cap \{x_n = 0\}$ be such that $\tau \cdot e > 0$.*

Then, for any given $\eta > 0$ we have

$$\partial_\tau u_0 \geq c_0(\tau \cdot e) > 0 \quad \text{in } \{x' \cdot e \geq \eta > 0\} \cap B_2$$

and

$$\partial_\tau u_0 \geq c_0(\tau \cdot e) > 0 \quad \text{in } \{|x_n| \geq \eta > 0\} \cap B_2,$$

with c_0 depending only on η and ellipticity constants.

Proof. Since $u_0(x) = u_0(x' \cdot e, x_n)$ it suffices to show the result in dimension $n = 2$. In that case, we have $F(D^2 u_0) = 0$ in $\mathbb{R}^2 \setminus \{x_1 \leq 0\}$, and satisfies $\partial_{x_1 x_1} u_0 \geq 0$, $\partial_{x_1} u_0 \geq 0$ in $\mathbb{R}^2$. Then, by the interior Harnack inequality, and using $\|u_0\|_{L^\infty(B_1)} = 1$, it follows that

$$\partial_{x_1} u_0 \geq c > 0 \quad \text{in } \{x_1 \geq \eta > 0\} \cap B_2$$

and

$$\partial_{x_1} u_0 \geq c > 0 \quad \text{in } \{|x_2| \geq \eta > 0\} \cap B_2,$$

as desired. □

We can now give the:

Proof of Proposition 6.1. Let $r > 0$ be as in the proof of Proposition 5.2, and

$$v(x) = \frac{u(rx)}{\|u\|_{L^\infty(B_r)}}.$$

Then, v satisfies

$$\begin{aligned} F(D^2 v) &= 0 \quad \text{in } B_2 \setminus \{x_n = 0\}, \\ \min(-F(D^2 v), v - \varphi_r) &= 0 \quad \text{on } B_2 \cap \{x_n = 0\}. \end{aligned}$$

Moreover, $\|\varphi_r\|_{C^2(B_1)} \leq Cr^\epsilon$.

Thus, the function

$$w = v - \varphi_r$$

solves $F(D^2w + D^2\varphi_r) = 0$ in $B_2 \cap \{x_n > 0\}$, and $\min(-F(D^2w), w) = 0$ on $B_2 \cap \{x_n = 0\}$. Therefore, any derivative $\partial_\tau w$, with $\tau \in S^{n-1} \cap \{x_n = 0\}$, satisfies

$$M^+(\partial_\tau w) \geq -Cr^\epsilon \quad \text{and} \quad M^-(\partial_\tau w) \leq Cr^\epsilon \quad \text{in} \quad B_2 \setminus \Omega^*,$$

where $\Omega^* := \{w = 0\} \cap \{x_n = 0\} \cap B_2$. Moreover, we have

$$\partial_\tau w = 0 \quad \text{on} \quad \Omega^*.$$

Now, notice that by Proposition 5.2, for any given $\delta > 0$ we may choose $r > 0$ small enough so that $|\partial_\tau w - \partial_\tau u_0| \leq \delta$, where u_0 is a global solution of (4.1)-(4.2)-(4.3). By Lemma 6.2, we find

$$\partial_\tau w \geq c_0(\tau \cdot e) - \delta \quad \text{in} \quad (\{x' \cdot e \geq \eta\} \cup \{|x_n| \geq \eta\}) \cap B_2.$$

Now, choosing δ small enough (depending on ℓ), this gives

$$\partial_\tau w \geq \tilde{c}_0 \quad \text{in} \quad (\{x' \cdot e \geq \eta\} \cup \{|x_n| \geq \eta\}) \cap B_2,$$

for all $\tau \in S^{n-1} \cap \{x_n = 0\}$ such that $\tau \cdot e \geq \ell/\sqrt{1+\ell^2}$. Finally, using Lemma 2.10 (applied to $\partial_\tau w$) we obtain

$$\partial_\tau w \geq 0 \quad \text{in} \quad B_1,$$

as desired. $\square$

7. THE REGULAR SET IS OPEN AND C^1

In this Section, we finally prove Theorem 1.1. By Proposition 6.1, we know that if x_0 is a regular point, then the free boundary is C^1 at x_0 . We next prove that the regular set is open, and this will yield Theorem 1.1.

In this section ϵ_0 denotes a fixed constant in $(0, 1 - \beta_2)$, where β_2 is “subsolution” exponent given by Proposition 2.7. For instance, we may fix $\epsilon_0 = \frac{1}{2}(1 - \beta_2)$, a constant depending only on the ellipticity constants (and thus universal).

Proposition 7.1. *Assume 0 is a regular free boundary point with exponent ϵ and modulus ν . Then, there is $e \in S^{n-1} \cap \{x_n = 0\}$ and there is $r > 0$ such that for any free boundary point $x_0 \in \partial\{u = \varphi\} \cap \{x_n = 0\} \cap B_r$ we have*

$$(u - \varphi)(x_0 + te) \geq ct^{2-\epsilon_0} \quad \text{for all} \quad t \in (0, r/2).$$

The constant $c > 0$ depends only on n, ϵ, ν , and ellipticity constants. In particular, every free boundary point in B_r is regular, with a uniform exponent $\epsilon = \epsilon_0/2$ and a uniform modulus $\tilde{\nu} = \tilde{\nu}(t) = ct^{\epsilon-\epsilon_0}$.

To prove Proposition 7.1, we need the following Lemma. Recall that x' denote points in $\mathbb{R}^{n-1}$ and the extension operators E^+ and E^- were defined in Proposition 2.6.

Lemma 7.2. *Let e be a unit vector in $\mathbb{R}^{n-1} \times \{0\}$, and $0 < \beta_1 < \frac{1}{2} < \beta_2 < 1$ the exponents in Corollary 2.9. Define*

$$\psi_{\text{sub}}(x') := e \cdot x' - \eta|x'| \left(1 - \frac{(e \cdot x')^2}{|x'|^2}\right)$$

$$\psi_{\text{super}}(x') := e \cdot x' + \eta |x'| \left(1 - \frac{(e \cdot x')^2}{|x'|^2} \right),$$

$$\Phi_{\text{sub}} := E^- \left[(\psi_{\text{sub}})_+^{\beta_2 + \gamma} \right] \quad \text{and} \quad \Phi_{\text{super}} := E^+ \left[(\psi_{\text{super}})_+^{\beta_1 - \gamma} \right].$$

For every $\gamma \in (0, \min\{|\beta_1 - 0|, |\beta_2 - 1|\})$ there is $\eta > 0$ such that two functions Φ_{sub} and Φ_{super} satisfy

$$\begin{cases} M^- \Phi_{\text{sub}} = 0 & \text{in } \{x_n > 0\} \\ \partial_{x_n} \Phi_{\text{sub}} \geq c_\gamma d^{\beta_2 + \gamma - 1} > 0 & \text{on } \{x_n = 0\} \cap \mathcal{C}_\eta^* \\ \Phi_{\text{sub}} = 0 & \text{on } \{x_n = 0\} \setminus \mathcal{C}_\eta^* \end{cases}$$

and

$$\begin{cases} M^+ \Phi_{\text{super}} = 0 & \text{in } \{x_n > 0\} \\ \partial_{x_n} \Phi_{\text{super}} \leq -c_\gamma d^{\beta_2 + \gamma - 1} < 0 & \text{on } \{x_n = 0\} \cap \mathcal{C}_{-\eta}^* \\ \Phi_{\text{super}} = 0 & \text{on } \{x_n = 0\} \setminus \mathcal{C}_{-\eta}^* \end{cases}$$

where $\mathcal{C}_{\pm\eta}^*$ is the cone

$$\mathcal{C}_{\pm\eta}^* := \left\{ (x', 0) \in \mathbb{R}^n : e \cdot \frac{x'}{|x'|} > \pm \eta \left(1 - \left(e \cdot \frac{x'}{|x'|} \right)^2 \right) \right\}, \quad (7.1)$$

and d is the distance to $\mathcal{C}_{\pm\eta}^*$. The constants c_γ and η depend only on γ , s , ellipticity constants, and dimension.

Proof of Lemma 7.2. We prove the statement for Φ_{sub} . The statement for Φ_{super} is proved similarly.

Let us denote $\psi = \psi_{\text{sub}}$ and $\Phi = \Phi_{\text{sub}}$. Note that Φ is the E^- extension of a homogeneous function of degree $\beta_2 + \gamma$ and thus by uniqueness of the extension (among functions with subcritical growth) it will be homogeneous with the same exponent.

By definition we have $M^- \Phi = 0$ in $\{x_n > 0\}$ and $\Phi = 0$ on $\{x_n = 0\} \setminus \mathcal{C}_\eta^*$ since $\psi < 0$ on that set.

We thus only need to check that, for $\eta > 0$ small enough

$$\partial_{x_n} \Phi \geq 0 \quad \text{on } \{x_n = 0\}$$

By homogeneity, it is enough to prove that $\partial_{x_n} \Phi \geq 0$ on points belonging to $e + \partial \mathcal{C}_\eta^*$, since all the positive dilations of this set with respect to the origin cover the interior of $\mathcal{C}_\eta^*$.

Let thus $P \in \partial \mathcal{C}_\eta^*$, that is,

$$e \cdot P = \eta \left(|P| - \frac{(e \cdot P)^2}{|P|} \right).$$

We note that —recall that both $P, e \in \{x_n = 0\}$

$$\begin{aligned}
\psi(P + e + x') &= e \cdot (P + e + x') - \eta \left(|P + e + x'| - \frac{(e \cdot (P + e + x'))^2}{|P + e + x'|} \right) \\
&= 1 + e \cdot x - \eta \left(|P + e + x| - |P| - \frac{(e \cdot (P + e + x))^2}{|P + e + x|} + \frac{(e \cdot P)^2}{|P|} \right) \\
&= 1 + e \cdot x' - \eta \psi_P(x') \\
\psi_P(x') &:= |P + e + x'| - |P| - \frac{(e \cdot (P + e + x'))^2}{|P + e + x'|} + \frac{(e \cdot P)^2}{|P|}.
\end{aligned}$$

Then we define

$$\begin{aligned}
\Phi_{P,\eta}(x) &:= \Phi(P + e + x) \\
&= E^- \left[(x', 0) \mapsto (1 + e \cdot x' - \eta \psi_P(x'))_+^{\beta_2 + \gamma} \right] (x),
\end{aligned}$$

where

Note that the functions ψ_P satisfy

$$\psi_P(0) = 0,$$

$$|\nabla \psi_P(x')| \leq C \quad \text{in } \mathbb{R}^n \setminus \{-P - e\},$$

and

$$|D^2 \psi_P(x')| \leq C \quad \text{for } x' \in B_{1/2}^*,$$

where C does not depend on P (recall that $|e| = 1$).

Then, the (traces of) the family $\Phi_{P,\eta}$ satisfy

$$\Phi_{P,\eta} \rightarrow (1 + e \cdot x')_+^{\beta_2 + \gamma} \quad \text{in } C^2(\overline{B_{1/2}^*})$$

as $\eta \searrow 0$, uniformly in P .

Moreover,

$$|\Phi_{P,\eta} - (1 + e \cdot x')_+^{\beta_2 + \gamma}| \leq (C\eta|x'|)^{\beta_2 + \gamma}$$

with C independent of P .

Thus, since $\beta_2 + \gamma < 1$, Lemma 2.8 implies

$$\partial_{x_n} \Phi_{P,\eta}(0) \rightarrow \partial_{x_n} E^- \left[(x', 0) \mapsto (1 + e \cdot x')_+^{s+\gamma} \right] (0) = c(s, \gamma, \lambda) > 0,$$

uniformly in P as $\eta \searrow 0$.

In particular one can chose $\eta = \eta(\gamma, \lambda, \Lambda)$ so that $\partial_{x_n} \Phi_{P,\eta}(0) \geq c(s, \gamma, \lambda) > 0$ for all $P \in \partial \mathcal{C}_\eta^*$ and the lemma is proved. $\square$

We can now show Proposition 7.1.

Proof of Proposition 7.1. We want to show that there is $e \in S^{n-1} \cap \{x_n = 0\}$ and there is $r > 0$ such that for any free boundary point $x_0 \in \partial\{u = \varphi\} \cap \{x_n = 0\} \cap B_r$ we have

$$(u - \varphi)(x_0 + te) \geq ct^{2-\epsilon_0} \quad \text{for all } t \in (0, r/2). \quad (7.2)$$

This will follow using the subsolutions of Proposition 7.2 and Lemma 2.3, from a inspection of the Proof of Proposition (6.1). Recall that in all the paper ϵ_0 denotes some constant in $(0, 1 - \beta_2)$.

Indeed, given $\eta > 0$ by Proposition (6.1) we find $r > 0$ such that, for every $x_0 \in \partial\{u = \varphi\} \cap \{x_n = 0\} \cap B_r$

$$u > \varphi \quad \text{on} \quad B_{2r}^* \cap (x_1 + \mathcal{C}_\eta). \quad (7.3)$$

Then, similarly as in the proof of Proposition (6.1) the function

$$w(x) = \frac{u(rx) - \varphi(rx)}{\|u\|_{L^\infty(B_r)}},$$

with $r > 0$ small satisfies

$$M^+(\partial_e w) \geq -\delta \quad \text{and} \quad M^-(\partial_e w) \leq \delta \quad \text{in} \quad B_2 \setminus \{x_n = 0, w = 0\},$$

where δ can be arbitrarily small provided that r is small enough.

Moreover, still as in the proof of Proposition (6.1), we have

$$\partial_e w \geq c_0 > 0 \quad \text{on} \quad B_1^* \cap \{x' \cdot e \geq 1/10\}. \quad (7.4)$$

Rescaling (7.3) we that hat, for every $x_0 \in \partial\{w = 0\} \cap B_1^*$

$$\{x_n = 0, w = 0\} \cap B_2^* \subset B_2^* \setminus (x_1 + \mathcal{C}_\eta)$$

Let us fix $\rho = 1/10$, $B^* = B_{1/4}^*(e/2)$, and $\gamma \in (\beta_2, 1)$ satisgying $\beta_2 + \gamma = 1 - \epsilon_0$. Let $\eta \in C^2(\overline{B_1})$ be some smooth “cutoff” function with $\eta = 1$ for $|x| \geq 1 - \rho$ and $\eta = 0$ in $B_{1/2}$. Let us call

$$C_1 := \sup_{B_1} M^+ \eta = \sup_{B_{1-\rho}} M^+ \eta > 0$$

Let ϕ be the subsolution of Lemma 2.3 with $\rho = 1/10$ and $B^* = B_{1/4}^*(e/2)$. Let $\Phi = \Phi_{\text{sub}} / \|\Phi_{\text{sub}}\|_{L^\infty(B_1)}$ the subsolution of Lemma 7.2 that vanishes in $\mathbb{R}^{n-1} \setminus \mathcal{C}_\eta^*$ and has homogeneity $\beta_2 + \gamma$.

Let us fix $x_0 \in \partial\{w = 0\} \cap B_1^*$.

We will show next that, for C large enough,

$$C\partial_e w - (x_n)^2 + 2\eta \geq 2C_1\phi + \Phi(\cdot - x_0) \quad \text{in} \quad B_1. \quad (7.5)$$

Let

$$v = \partial_e w - (x_n)^2 + 2\eta - 2C_1\phi - \Phi(\cdot - x_1).$$

On on hand, let us show that $v \geq 0$ on ∂B_1 . Indeed, we have (r is large) we have $\partial_e w \geq 0$ in B_1 . Also, $\eta = 1$ for for $|x| \geq 1 - \rho$ and thus $\eta - |x|^2 = 0$ on ∂B_1 . Moreover, recall that $\phi = 0$ on ∂B_1 and, since $0 \leq \Phi \leq 1$ in B_1 , $\eta - \Phi \geq 0$ on ∂B_1 .

On the other hand, let us show that

$$M^- v \leq 0 \quad \text{in} \quad (B_1 \setminus B^*) \cup (x_0 + \mathcal{C}_\eta^*).$$

Indeed, we have

$$\begin{aligned}
M^-v &= M^-(C\partial_e w - (x_n)_+^2 + 2\eta - 2C_1\phi - \Phi) \\
&\leq CM^-(\partial_e w) - 2\lambda + 2 \sup_{B_{1-\rho}} M^+\eta - 2C_1M^-\phi + M^+\Phi(\cdot - x_0) \\
&\leq C\delta - 2\lambda + 2C_1\chi_{B_{1-\rho}} - 2C_1\chi_{B_{1-\rho}} + M^+\Phi(\cdot - x_0) \\
&\leq C\delta - 2\lambda \\
&\leq 0
\end{aligned}$$

in $(B_1 \setminus B^*) \cup (x_1 + \mathcal{C}_\eta^*)$ provided that $C\delta - 2n\lambda \leq 0$.

That $v \geq 0$ in $B_1^* \setminus (x_0 + \mathcal{C}_\eta^*)$ is now a consequence of (7.3) which implies that $w = (x_n)^2 = \phi = \Phi = 0$ on that set. Last, recalling (7.4) we see that $v \geq 0$ in B^* can be guaranteed by choosing C large (depending only on c_0 and universal constants).

Thus, choosing first C large and then δ small enough so that $C\delta - 2n\lambda \leq 0$, and using the maximum principle, we prove $v \geq 0$ in B_1 and thus that

$$C\partial_e w \geq \Phi(\cdot - x_0) = (\psi_{\text{sub}}(\cdot - x_1))_+^{\beta_2+\gamma} \quad \text{on } B_{1/2}^*,$$

where ψ_{sub} was defined in Lemma 7.2.

After rescaling and noting that $\psi_{\text{sub}}(te) = t$, this implies that

$$\partial_e w(te) \geq ct^{\beta_2+\gamma} = ct^{1-\epsilon_0} > 0 \quad \text{for } t \in (0, r/2).$$

Thus, (7.2) follows integrating with respect to t (note that $w(0) = \partial_e(0) = 0$). $\square$

Finally, as a consequence of the previous results, we give the:

Proof of Theorem 1.1. By Proposition 7.1, the set of regular points is open, and (i) holds at all such points. Moreover, still by Proposition 7.1, given any free boundary point x_0 , there is a ball $B_r(x_0)$ in which all free boundary points are regular, with a common modulus of continuity ν . Thus, by Proposition 6.1, the free boundary is C^1 at each of these points, with a uniform modulus of continuity (that depends on x_0). Thus, the free boundary is locally a C^1 graph in $B_r(x_0)$. $\square$

When the ellipticity constants λ and Λ are close to 1, we establish the following.

Corollary 7.3. *Let F be as in (1.4), and u be any solution of (1.2), with $\varphi \in C^{1,1}$. Then, for any small $\delta > 0$ we have*

$$u \in C^{1, \frac{1}{2}-\delta}(B_{1/2}) \quad \text{whenever} \quad |\Lambda - 1| + |\lambda - 1| \leq \delta/C_0.$$

The constant C_0 is universal. Furthermore, under such assumption on the ellipticity constants, we have

$$\|u\|_{C^{1, \frac{1}{2}-\delta}(B_{1/2})} \leq C(\|u\|_{L^\infty(B_1)} + \|\varphi\|_{C^{1,1}(B_1 \cap \{x_n=0\})},$$

with C depending only on n , λ and Λ .

Proof. The proof is by contradiction, using the result in Corollary 4.4.

Dividing by a constant if necessary, we assume $\|u\|_{L^\infty(B_1)} + \|\varphi\|_{C^{1,1}(B_1 \cap \{x_n=0\})} \leq 1$. We first claim that, for every free boundary point $x_0 \in B_{1/2} \cap \partial\{u = \varphi\}$, we have

$$|u(x) - u(x_0) - \nabla u(x_0) \cdot (x - x_0)| \leq C|x - x_0|^{\frac{3}{2}-\delta}, \quad (7.6)$$

with C depending only on n and λ, Λ .

Let us prove (7.6) by contradiction. Indeed, assume there are sequences of operators F_k as in (1.4), obstacles φ_k satisfying $\|\varphi_k\|_{C^{1,1}(B_1 \cap \{x_n=0\})} \leq 1$, solutions u_k to (1.2) with $\|u_k\|_{L^\infty(B_1)} \leq 1$, and free boundary points $x_k \in B_{1/2}$, such that

$$|u_k(x) - \nabla u_k(x_k)| \geq k|x - x_k|^{\frac{3}{2}-\delta},$$

for all $k \geq 1$. By the $C^{1,\alpha}$ estimates in [Fer16], we know that $|\nabla u_k(x_0)| \leq C$, so that after subtracting a linear function we may assume $u_k(x_k) = 0$ and $\nabla u_k(x_k) = 0$. Moreover, after a translation we may assume for simplicity that $x_k = 0$.

Then, defining

$$\theta(\rho) = \sup_{\rho \leq r \leq 1} \sup_k r^{\delta-\frac{3}{2}} \|u_k\|_{L^\infty(B_r)},$$

and by the exact same argument in Lemma 5.4, we find a sequence $r_k \rightarrow 0$ for which

$$w_k(x) = \frac{u_k(r_k x)}{\|u_k\|_{L^\infty(B_{r_k})}}$$

satisfies

$$|w_k(x)| \leq C(1 + |x|^{\frac{3}{2}-\delta}) \quad \text{in } B_{1/r_k},$$

$\|w_k\|_{L^\infty(B_1)} = 1$, $w_k(0) = 0$, $\nabla w_k(0) = 0$, and

$$\begin{aligned} F_k(D^2 w_k) &= 0 \quad \text{in } B_{1/r_k} \setminus \{x_n = 0\} \\ \min(-F_k(D^2 w_k), w_k - \varphi_k) &= 0 \quad \text{on } B_{1/r_k} \cap \{x_n = 0\}, \end{aligned}$$

where

$$\|\varphi_k\|_{C^{1,1}(B_R)} = \frac{\|\varphi\|_{B_{Rr_k}}}{\|u_k\|_{L^\infty(B_{r_k})}} \leq \frac{CR^2(r_k)^2}{(r_k)^{\frac{3}{2}-\delta}} = CR^2(r_k)^{\frac{1}{2}+\delta}$$

converges to 0 for every fixed R as $k \rightarrow \infty$.

Thus, by $C^{1,\alpha}$ estimates, up to a subsequence the operators F_k converge to an operator F as in (1.4), and the functions w_k converge locally uniformly to a function w satisfying

$$|w(x)| \leq C(1 + |x|^{\frac{3}{2}-\delta}) \quad \text{in } \mathbb{R}^n,$$

$\|w\|_{L^\infty(B_1)} = 1$, $w(0) = 0$, $\nabla w(0) = 0$, and

$$\begin{aligned} F(D^2 w) &= 0 \quad \text{in } \mathbb{R}^n \setminus \{x_n = 0\} \\ \min(-F(D^2 w), w) &= 0 \quad \text{on } \mathbb{R}^n \cap \{x_n = 0\}. \end{aligned}$$

By Corollary 4.4, we get $w \equiv 0$, a contradiction. Thus, (7.6) is proved.

Finally, combining (7.6) with interior regularity estimates, the result follows exactly as in the proof of [Fer16, Theorem 1.1]. $\square$

REFERENCES

- [ACS08] I. Athanassopoulos, L. Caffarelli, S. Salsa, *The structure of the free boundary for lower dimensional obstacle problems*, Amer. J. Math. 130 (2008) 485-498.
- [BFR15] B. Barrios, A. Figalli, X. Ros-Oton, *Global regularity for the free boundary in the obstacle problem for the fractional Laplacian*, preprint arXiv (June 2015).
- [BB94] R. Bass, K. Burdzy, *The boundary Harnack principle for non-divergence form elliptic operators*, J. Lond. Math. Soc. 50 (1994), 157-169.
- [Caf77] L. Caffarelli, *The regularity of free boundaries in higher dimensions*, Acta Math. 139 (1977), 155-184.
- [Caf79] L. Caffarelli, *Further regularity for the Signorini problem*, Comm. Partial Differential Equations 4 (1979) 1067-1075.
- [Caf80] L. Caffarelli, *Compactness methods in free boundary problems*, Comm. Partial Differential Equations 5 (1980), 427-448.
- [Caf98] L. Caffarelli, *The obstacle problem revisited*, J. Fourier Anal. Appl. 4 (1998), 383-402.
- [CC95] L. Caffarelli, X. Cabré, *Fully Nonlinear Elliptic Equations*, Amer. Math. Soc. Colloq. Publ., vol. 43, American Mathematical Society, Providence, RI, 1995.
- [CRS16] L. Caffarelli, X. Ros-Oton, J. Serra, *Obstacle problems for integro-differential operators: regularity of solutions and free boundaries*, preprint arXiv (Jan. 16).
- [CSS08] L. Caffarelli, S. Salsa, L. Silvestre, *Regularity estimates for the solution and the free boundary of the obstacle problem for the fractional Laplacian*, Invent. Math. 171 (2008), 425-461.
- [DS14] D. De Silva, O. Savin, *Boundary Harnack estimates in slit domains and applications to thin free boundary problems*, Rev. Mat. Iberoam., to appear.
- [Fer16] X. Fernández-Real, *$C^{1,\alpha}$ estimates for the fully nonlinear Signorini problem*, preprint arXiv (Mar. 16).
- [Fre77] J. Frehse, *On Signorini's problem and variational problems with thin obstacles*, Ann. Scuola Norm. Sup. Pisa 4 (1977), 343-362.
- [GP09] N. Garofalo, A. Petrosyan, *Some new monotonicity formulas and the singular set in the lower dimensional obstacle problem*, Invent. Math. 177 (2009), 415-461.
- [GPS15] N. Garofalo, A. Petrosyan, M. Smit Vega Garcia, *An isoperimetric inequality approach to the regularity of the free boundary in the Signorini problem with variable coefficients*, J. Math. Pures Appl., to appear.
- [Kin81] D. Kinderlehrer, *The smoothness of the solution to the boundary obstacle problem*, J. Math. Pures Appl. 60 (1981), 193-212.
- [KPS15] H. Koch, A. Petrosyan, W. Shi, *Higher regularity of the free boundary in the elliptic Signorini problem*, Nonlinear Anal. 126 (2015), 3-44.
- [KRS15] H. Koch, A. Rüland, W. Shi, *The variable coefficient thin obstacle problem: optimal regularity and regularity of the regular free boundary*, preprint arXiv (Apr. 2015).
- [Lew68] H. Lewy, *On a variational problem with inequalities on the boundary*, Indiana J. Math. 17 (1968), 861-884.
- [MS08] E. Milakis, L. Silvestre, *Regularity for the nonlinear Signorini problem*, Adv. Math. 217 (2008), 1301-1312.
- [RS15] X. Ros-Oton, J. Serra, *Boundary regularity estimates for nonlocal elliptic equations in C^1 and $C^{1,\alpha}$ domains*, preprint arXiv (Dec. 15).

THE UNIVERSITY OF TEXAS AT AUSTIN, DEPARTMENT OF MATHEMATICS, 2515 SPEEDWAY,
AUSTIN, TX 78751, USA

E-mail address: `ros.oton@math.utexas.edu`

WEIERSTRASS INSTITUT FÜR ANGEWANDTE ANALYSIS UND STOCHASTIK, MOHRENSTRASSE
39, 10117 BERLIN, GERMANY

E-mail address: `joaquim.serra@upc.edu`